

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
19 December 2002 (19.12.2002)

PCT

(10) International Publication Number  
**WO 02/101049 A2**

(51) International Patent Classification<sup>7</sup>: **C12N 15/24**,  
C07K 14/54, A61K 38/20, A61P 31/12, 35/00

(21) International Application Number: **PCT/IB02/02344**

(22) International Filing Date: **8 March 2002 (08.03.2002)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
60/274,327 **8 March 2001 (08.03.2001)** **US**

(71) Applicant: **ARES TRADING S.A. [CH/CH];** Le  
Château, CH-2028 Vaumarcus (CH).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors: **DINARELLO, Charles**; 333 15th Street, Boulder, CO 80302 (US). **KIM, Soo-Hyun**; 860 South Oneida Street, #C-112, Denver, CO 80224 (US).

**Published:**

— *without international search report and to be republished upon receipt of that report*

(74) Agent: **SERONO INTERNATIONAL S.A.**; Intellectual Property, 12, chemin des Aulx, CH-1228 Plan-les-Ouates (CH).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



**WO 02/101049 A2**

(54) Title: **INTERLEUKIN-18 MUTANTS, THEIR PRODUCTION AND USE**

(57) Abstract: The invention provides mutants of IL-18 with lower affinity to IL-18BP than the wild type IL-18 molecule.

5                   **INTERLEUKIN-18 MUTANTS, THEIR PRODUCTION AND USE**  
**FIELD OF THE INVENTION**

                  The present invention relates to IL-18 mutants having enhanced biological  
10   activity with respect to the wild type protein.

**BACKGROUND OF THE INVENTION**

                  In 1989, an endotoxin-induced serum activity that induced interferon- $\gamma$  (IFN- $\gamma$ )  
obtained from mouse spleen cells was described (Nakamura et al., 1989). This serum  
15   activity functioned not as a direct inducer of IFN- $\gamma$  but rather as a co-stimulant  
together with IL-2, IFN- $\gamma$ , TNF or mitogens. An attempt to purify the activity from  
post-endotoxin mouse serum revealed an apparently homogeneous 50-55 kDa protein  
(Nakamura et al., 1993). Since other cytokines can act as co-stimulants for IFN- $\gamma$   
production, the failure of neutralizing antibodies to IL-1, IL-4, IL-5, IL-6, or TNF to  
20   neutralize the serum activity suggested it was a distinct factor. In 1995, the same  
scientists demonstrated that the endotoxin-induced co-stimulant for IFN- $\gamma$  production  
was present in extracts of livers from mice preconditioned with *P. acnes* (Okamura et  
al., 1995). In this model, the hepatic macrophage population (Kupffer cells) expand  
and in these mice, a low dose of bacterial lipopolysaccharide (LPS), which in non-  
25   preconditioned mice is not lethal, becomes lethal. The factor, named IFN- $\gamma$  -inducing  
factor (IGIF) and later designated interleukin-18 (IL-18), was purified to homogeneity  
from 1,200 grams of *P. acnes*-treated mouse livers. Degenerate oligonucleotides  
derived from amino acid sequences of purified IL-18 were used to clone a murine IL-  
18 cDNA (Okamura et al. 1995). Messenger RNAs for IL-18 and interleukin-12 (IL-  
30   12) are readily detected in activated macrophages. IL-18 does not induce IFN- $\gamma$  by  
itself, but functions primarily as a co-stimulant with mitogens or IL-12. The human  
cDNA sequence for IL-18 was reported in 1996 (Figure 1A SEQ ID NO:1).

Interleukin IL-18 shares structural features with the IL-1 family of proteins (Nakamura et al., 1993; Okamura et al., 1995; Ushio et al., 1996; Bazan et al., 1996). Unlike most other cytokines, which exhibit a four-helix bundle structure, IL-18 and IL-1 $\beta$  have an all  $\beta$ -pleated sheet structure (Tsutsui et al., 1996). Similarly to IL-1 $\beta$ , IL-18 is synthesised as a biologically inactive precursor (proIL-18), lacking a signal peptide (Ushio et al., 1996). The IL-1 $\beta$  and IL-18 precursors are cleaved by caspase 1 (IL-1 $\beta$ -converting enzyme, or ICE), which cleaves the precursors after an aspartic acid residue in the P1 position. The resulting mature cytokines are readily released from the cell (Ghayur et al., 1997 and Gu et al., 1997).

IL-18 is a co-stimulant for cytokine production (IFN- $\gamma$ , IL-2 and granulocyte-macrophage colony stimulating factor) by T helper type I (Th1) cells (Kohn et al., 1997) and also a co-stimulant for FAS ligand-mediated cytotoxicity of murine natural killer cell clones (Tsutsui et al., 1996).

Th1 lymphocytes are involved in the immune responses against tumors (Seki et al., 2000). Th1 responses include the secretion of the cytokines IL-2, IL-12, IL-18 and IFN- $\gamma$ , as well as the generation of specific cytotoxic T lymphocytes recognizing specific tumor antigens. The Th1 response is also a vital arm of host defence against many microorganisms. However, the Th1 response can also be associated with non-desirable effects such as the development of several autoimmune diseases, inflammation and organ transplant rejection.

Attempts to express the mature form of IL-18 in *E. coli* using a vector encoding the mature protein did not provide a fully active cytokine. An efficient expression system for the generation of the fully biologically active human IL-18 has been developed for therapeutic uses, e.g. in malignancies, or any condition where IFN- $\gamma$  induction is desired (WO 00/61768). In this system, the IL-18 precursor caspase-1 cleavage site has been changed to a factor Xa site (ICE/Xa), and a vector encoding IL-18 ICE/Xa precursor was used for transformation of *E. coli*. Following expression of this IL-18 precursor in *E. coli* the mature IL-18 was generated by factor Xa cleavage *in vitro*. This mature IL-18 generated by factor Xa cleavage was fully active.

Cytokine binding proteins (soluble cytokine receptors) are usually the extracellular ligand binding domains of their respective cell surface cytokine receptors. They are produced either by alternative splicing or by proteolytic cleavage of the cell surface receptor. These soluble receptors have been described in the past, for example, the soluble receptors of IL-6 and IFN- $\gamma$  (Novick et al., 1989), TNF (Engelmann et al., 1989; Engelmann et al., 1990), IL-1 and IL-4 (Maliszewski et al., 1990), IFN- $\alpha/\beta$  (Novick et al., 1994; Novick et al. 1992). One cytokine-binding protein, named osteoprotegerin (OPG, also known as osteoclast inhibitory factor-OCIF), a member of the TNFR/Fas family, appears to be the first example of a soluble receptor that exists only as a secreted protein (Anderson et al., 1997; Simonet et al., 1997; Yasuda et al., 1998).

An IL-18 binding protein (IL-18BP) was affinity purified, on an IL-18 column, from urine (Novick et al., 1999). IL-18BP abolishes IL-18 induction of IFN- $\gamma$  and of IL-8, activation of NF- $\kappa$ B *in vitro* and induction of IFN- *in vivo*. IL-18BP is a soluble circulating protein which is constitutively expressed in the spleen, and belongs to the immunoglobulin superfamily. The most abundant IL-18BP isoform, the spliced variant isoform a, exhibits a high affinity for IL-18 with a rapid on-rate and a slow off-rate, and a dissociation constant (Kd) of approximately 400 pM (Kim et al., 1999).

The residues involved in the interaction of IL-18 with IL-18BP have been described through the use of computer modelling (Kim et al., 1999) and based on the interaction of IL-1 with the IL1R type I (Vigers et al., 1997). In the model for IL-18 binding to the IL-18BP, the Glu residue at position 42 and the Lys residue at position 89 of IL-18 have been proposed to bind to Lys-130 and Glu-114 in IL-18BP, respectively (Kim et al., 1999).

IL-18 is constitutively present in many cells (Puren et al., 1999) and circulates in healthy humans (Urushihara et al., 2000). The high affinity of IL-1BP to IL-18 as well as the high concentration of IL-18BP found in the circulation (20 fold molar excess over IL-18), represents a unique situation in cytokine biology. Therefore, most, if not all, of the IL-18 molecules in the circulation is bound to the IL-18BP.

The circulating IL-18BP which competes with cell surface receptors for IL-18, may act as a natural anti-inflammatory and an immunosuppressive molecule.

Viral agents encode IL-18BP like proteins, for example, *M. contagiosum* viral proteins MC53 and MC54 share a significant homology to mammalian IL-18BP  
5 (Novick et al. 1999). *M. contagiosum* proteins MC53 and MC54 possess the ability to bind and neutralize human IL-18 in a fashion similar to that of IL-18B (Xiang and Moss, 1999). The ectromelia poxvirus p13 protein, which is homologous to IL-18BP, binds human IL-18 and inhibits its activity in vitro. Mice infected with a p13 deletion mutant virus exhibited decreased levels of infectivity (Born et al., 2000). Therefore  
10 infectivity degree seems to correlate with the presence of IL-18BP.

The high levels of circulating IL-18BP may represent a natural defence against a runaway Th1 response to infection and development of autoimmune diseases. However, IL-18 contributes to the Th1 response which is important in host defence against tumors. Therefore, IL-18BP may bring about failure of the host to develop  
15 cytotoxic T cells directed against tumor cells. Indeed, there is evidence that IL-18 promotes host defence against tumors in mice. For example, in syngeneic mice, murine mammary carcinoma cells expressing murine IL-12 or murine IL-18 were less tumorigenic and formed tumors more slowly than did control non-expressing cells (Coughlin et al., 1998). Antibody neutralisation studies revealed that the antitumor  
20 effects required IFN- $\gamma$ . In another study, systemically administration of IL-18 to experimental animals in combination with B16 melanoma expressing B7-1 (CD80) resulted in dramatic suppression of melanoma formation, tumor growth, and a significant improvement in survival (Cho et al., 2000).

Cytokines are used as adjuvant to increase the effectivity of immunotherapy in  
25 cancer. For example, IL-2 is administered for renal cell carcinoma or melanoma (Gollob et al., 2000). Often, one important consequence of the treatment with cytokines is severe systemic toxicity profiles. Using cytokines, expressed by the patient's own tumor or dendritic cells, is a logical solution to the problem. Yet, if IL-18 will to be used locally, as adjuvant in tumor immunotherapy, the ability of the  
30 constitutive levels of IL-18BP to neutralize IL-18 in the local environment would still be exerted and consequently its effectivity is greatly diminished.

The use of non-myeloablative allogeneic transplants, the so-called mini transplants, to treat leukaemia and solid tumors is increasingly successful in inducing graft-versus-leukaemia and graft-versus-tumor reactions (Slavin S., 2000; Slavin et al., 2000). Two studies that used either allogeneic peripheral blood stem cells (Childs et al. 2000) or dendritic cells (Kugler et al., 2000) to treat patients with metastatic renal cell carcinoma met a remarkable success. Although these studies need to be extended and confirmed, the concept that an ongoing graft-versus-tumor reaction is exploitable for immunotherapy in cancer is gaining acceptance (Slavin, 2000). Since IL-18 appears to be involved in these successful therapeutic approaches, a further improvement may be achieved if the neutralizing effect of IL-18BP can be abolished.

Mutants of IL-18 (IFN- $\gamma$ , inducing factor) are described in EP0845530. The described IL-18 mutants are molecules in which 1, 2, 3 or all 4 cysteine residues in IL-18 (Figure 1B) were replaced by serine or alanine residues. These mutants contained an intact consensus sequence (Figure 1B). All the isolated mutants exhibit higher stability than wild type IL-18. The degree of stability of the mutants is directly proportional to the number of Cys residues replaced in the molecule. EP0845530 is silent on the ability of IL-18BP to neutralize these mutants.

The generation and therapeutic use of fully active IL-18 mutants unable to bind or bind with low affinity to IL-18BP, is therefore highly advantageous.

## SUMMARY OF THE INVENTION

The present invention relates to an IL-18 mutant polypeptide, comprising mutations in one or more amino acid residues which are involved in its interaction with IL-18 binding protein. More specifically, the mutations are substitutions, preferably non-conservative, additions or deletions. The residues mutated in said polypeptide may be selected from Glu-42, Ile-85, Met-87, Lys-89, Met-96, Asp-130, Lys-132, Pro-143, Met-149, and Leu-189, preferably, Glu-42 and Lys-89.

In one embodiment, the Glu-42 or Lys-89 or both Glu-42 and Lys-89 are replaced with a non-polar amino acid, preferably alanine.

In addition, the invention provides for a DNA encoding said polypeptide, preferably encoding a polypeptide of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7 or SEQ ID NO:8.

In one embodiment, DNA encoding SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8 is fused to a signal peptide, preferably that of the hGH. Moreover, the invention also provides a vector comprising said DNA capable of expressing the polypeptide encoded by said DNA in an appropriated host cell, e.g. a prokaryotic or eukaryotic host cell.

In addition, the present invention provides for pharmaceutical compositions comprising said a polypeptide for the treatment of diseases which are prevented or alleviated by Th1 responses, preferably for treatment of viral disease or cancer.

#### BRIEF DESCRIPTION OF THE FIGURES

Figure 1A shows the nucleotide sequence encoding the WT IL-18 precursor and the location of primers used to construct the different mutated IL-18 proteins. The broad arrow indicates where the mature IL-18 protein coding sequence begins.

Figure 1B shows the amino acid sequence of the mature IL-18. The consensus sequence among different species of IL-18 are enclosed in the white boxes. The cystidines are underlined. The dark boxes show the residues mutated in IL-18 according to this invention.

Figure 2 shows the schematic representation of the IL-18 mutants according to the invention. The His-6 indicates the location of the six histidines fused in the N terminus of the IL-18 precursor propiece. The arrow indicates the ICE cleavage site as replaced by the factor Xa cleavage site (x). WT indicates the wild-type mature IL-18. E42A indicates Glu-42 to Ala mutation, K89A indicates Lys-89 to Ala mutation and E42A/K89A indicates double mutation. On the basis of precursor /x/ WT, three IL-18 mutants (E42A, K89A and E42A+K89A) were generated by two step PCR.

Figure 3 A shows the induction of IFN- in NKO cells by IL-18 WT and mutant protein at concentrations shown under the x-axis in Fig. 3B and in the presence of IL-12 (0.5 ng/ml)

Figure 3 B shows the induction of IFN- in PBMCs cells by IL-18 WT and mutant protein at concentrations shown under the x-axis and in the presence of IL-12 (1.0 ng/ml).

Figure 4 A shows the effect of IL-18BP in IFN- induction by human IL-18 WT and mutant protein in NK0 cells. Mutants and WT IL-18 (30ng/ml) were preincubated with IL-18BP at the concentrations indicated under the x-axis (of Fig. 3B) for 1 h at room temperature and added to NK0 cells stimulated with IL-12 ( 0.5 mg/ml ).

Figure 4 B shows the effect of IL-18BP in IFN- induction by human IL-18 WT and mutant protein in PBMCs cells. Mutants and WT IL-18 (30ng/ml) were preincubated with IL-18BP at the concentrations, indicated under the x-axis for 1 h at room temperature and added to PBMCs cells stimulated with IL-12 (1.0 mg/ml).

Figure 5 shows the induction of IL-8 by IL-18 WT and mutant protein. PBMCs were incubated with IL-18 WT or mutant (30 ng/ml). Polymyxin B (1 g/ml) was mixed with IL-18 for 30 min before being added to the PBMCs. After 24 h the supernatants was removed and assayed for IL-18 concentration by ECL (example 9). One out of three experiments is shown.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to an IL-18 mutant or its active fragment, or a mutein, or any other protein or peptide derivative thereof (IL-18M), which is less susceptible to neutralization by IL18BP as compared to the wild type (IL-18 WT). More specifically, one or more amino acids of the IL-18 WT, preferably no more than 30, more preferably up to 10 amino acids, may be replaced with other amino acids, or eliminated, or one or more amino acids may be added, preferably no more than 30, more preferably up to 10 amino acid in order to generate an active IL-18 mutant which is less susceptible to neutralization by IL-18BP. Amino acids may be replaced by different amino acids the substitutions preferably being non-conservative substitutions. More specifically, said mutations could be targeted to residues predicted to be involved in binding to IL-18BP such as Glu-42, Ile-85, Met-87, Lys-



89, Met-96, Asp-130, Lys-132, Pro-143, Met-149, and Leu-189, more preferably Glu-42 and/or Lys-89 (Kim et al., 1999).

IL-18M may be produced, in eukaryotic or eukaryotic expression systems, intracellularly, periplasmic or may be secreted into the medium. The produced IL-18M may be recovered in soluble or insoluble form (inclusion bodies).

A vector comprising the precursor IL-18M cDNA may be used for expression of correct assembled precursor IL-18M in prokaryotic systems. Subsequently the mature fully active protein can be generated after cleavage by ICE *in vitro*. A sequence encoding the specific cleavage site for a protease, preferably factor Xa, can replace the ICE sequence in the precursor of IL-18M.

An expression vector encoding an effective signal peptide, preferably the human growth hormone signal peptide, fused to the cDNA of the mature IL-18M may be used for eukaryotic expression and secretion.

The parental IL-18 cDNA used for the mutant construction can be selected from mouse or human species.

IL-18M can be epitope tagged, preferably with histidine, for convenient purification. The recombinant IL-18M may be purified by conventional or affinity methods. The amount of IL-18M produced may be monitored by a specific ELISA.

IL-18M can be used in a pharmaceutical formulation for treatment of diseases which are prevented or reduced by Th1 responses, more specifically by IL-18 treatment e.g. microorganism infections and cancer. The advantage of using the mutant instead of the wild type version of the protein, resides in its resistance to IL-18BP neutralization.

More specifically, IL-18M may be administered systemically or locally as an adjuvant for tumor antigens in tumor immunotherapy.

Tumor cells derived from a patient can be isolated and genetically modified to secrete IL-18M and re-grafted to the same patient for local vaccination (Coughlin et al., 1998). Fusion of the modified tumor cells expressing IL-18M to allogeneic dendritic cells (antigen presenting cells) can be carried out to further increase the tumor antigenicity and consequently, the anti-tumor response.

IL-18M can be administered as an adjuvant in DNA vaccination (Tuting et al., 1998). In manner similar to that reported for the use of the cytokines IL-12 and IFN-. In this case, transdermal tumor antigen vaccination can be performed by using a gene gun. This may result in transfection of resident dendritic cells in the skin with DNA  
5 encoding both tumor antigen and IL-18M. Alternatively, the dendritic cells can be engineered *ex vivo* followed by an adoptive transfer.

IL-18M can be used as an adjuvant in graft versus tumor therapy. Allogeneic stem cells can be used to transplant cancer patients. To increase the graft versus tumor rejection induced by transplantation of the allogeneic cells, IL-18M may be  
10 administered systemically or locally, preferably by IL-18M expression in genetically modified syngeneic dendritic cells or in tumor cells taken from the patient.

It is to be understood that while the invention has been described in conjunction with the preferred specific embodiments thereof that the foregoing description as well as the examples that follow are intended to illustrate and not limit  
15 the scope of the invention. Other aspects, advantages and modification within the scope of the invention will be apparent to those skilled in the art to which the invention pertains.

## EXAMPLES

### 20 Example 1

**Construction of vectors for the expression of WT proIL-18 Histidine tag fusion protein for cleavage by Factor Xa.** In order to generate the correct assembled IL-18 in *E.coli*, the ICE cleavage site in the precursor of IL-18 was replaced by a Xa cleavage site. Subsequently in vitro, cleavage of the IL-18 precursor  
25 by Xa generates therefore the active protein (WO 00/61768).

The cDNA sequence encoding the human IL-18 precursor (proIL18, gene bank accession number D49950, Figure 1) used for generating the expression plasmid was isolated as described (Ghayur et al., 1997).

The replacement of the ICE cleavage site with a Xa cleavage site was achieved  
30 by using 2 PCR reactions (see primers used in Figure 1). PCR reaction 1: The propiece of IL-18 cDNA was generated by using the sense primer (Pr 1) containing

the EcoRI site located upstream of the ORF,

5'-ATATGAATTCATGGCTGCTGAACCAGTAG, (SEQ ID NO: 11) and a reverse primer (Pr 2) designed for the ICE site (33-LESD-36) in which 6 nucleotides have been changed to encode the factor Xa site (33-IEGR-36),

- 5 5'-AAAGTAACGTCCTTCGATGTTTTTC (SEQ ID NO: 12). The amplified DNA fragment encoding the mature IL-18 was generated by using the sense primer (Pr 3) which is complementary to Pr2,

5'-GAAAACATCGAAGGACGTTACTTT, (SEQ ID NO: 13) and the reverse primer (Pr 4) containing the BamHI site downstream of the coding sequence of IL18

- 10 5'-ATATGGATCCTAGTCTTCGTTTTGAACAGTG (SEQ ID NO: 14). The propiece 108 bp and mature 474 bp IL-18 DNAs were resolved by electrophoresis in 1% agarose, and eluted by a gel extraction system (GIBCO/ BRL).

- PCR reaction 2: The two DNA fragments obtained in the PCR reaction 1 were mixed at a 1:1 ratio and used together with primers Pr 1 and Pr 4 to generate a  
15 complete human IL-18 cDNA in which the ICE site is replaced by the factor Xa site (ICE/Xa).

- The pro IL-18 (ICE/Xa) cDNA was ligated into the BlueScript plasmid (Stratagene) by EcoRI and BamHI (GIBCO/BRL) restriction sites. This plasmid served for sequence confirmation. The predicted amino acid sequence encoded is  
20 shown in SEQ ID NO:2. For *E. coli* expression, the IL-18 DNA insert was re-ligated into the pPROEX HTa vector (GIBCO/BRL) with the use of EcoRI and XbaI sites (originating in BlueScript). In this vector, the resulting protein is N-terminal fused to a histidine tag.

## 25 **Example 2**

- Construction of the E42A, K89A and E42A/K89A mutants.** Mutations in IL-18 were created in residues predicted to be important for the binding to the inhibitor IL-18BP( Kim et al. 2000). Three mutants: E42A, K89A, and F-42A/K89A, were generated. The mutations were achieved by two PCR reactions, as described in  
30 example 1, using the primers and templates described below (the primers are shown in Figure 2).

E42A mutant

PCR reaction 1- The pair of primes used for preparing the mutant E42A were:  
pair A- Pr 1 (example 1) and the reverse primer (Pr 5) 5'- TAA TTT AGA TGC  
AAG CTT GCC (SEQ ID NO:15) encoding alanine instead of glutamic acid (E 42),  
5 and Pair B the sense primer (Pr 6) 5'-GGC AAG CTT GCA TCT AAA TTA (SEQ  
ID NO:16) encoding Alanine in exchange for Glutamic acid (GAA to GCA) and the  
reverse primer Pr 4 (example 1), using pro IL-18 (ICE/Xa) as a template in the PCR  
reaction.

PCR reaction 2: The two DNA fragments obtained in PCR reactions 1 were  
10 used as templates for the second PCR reaction using primers Pr1 and Pr 4.

K89A mutant

PCR reaction 1: The pair of primes used for preparing the mutant K89A were  
pair A Pr 1(example 1) and the reverse primer (Pr 7) 5'-CTG GCT ATC TGC ATA  
CAT ACT (SEQ ID NO:17) encoding alanine instead of lysine (K 89), and pair B the  
15 sense primer (Pr 8) 5'-AGT ATG TAT GCA GAT AGC CAG (SEQ ID NO:18)  
encoding alanine instead of Lysine (AAA to GCA) with the reverse primer Pr 4  
(Example 1) using pro IL-18 (ICE/Xa) as the template for the first PCR reaction.

PCR reaction 2: The two DNA fragments obtained in the PCR reaction 1 were  
used as templates for the second PCR reaction using primers Pr1 and Pr 4 .

20 E42A/K89A mutant

For the double mutation E42A/K89A, the primer used were the same as for the  
preparation of the F-42A mutation and mutant K89A cDNA was used as the template  
in the reaction.

Each of the three IL-18 mutated genes were ligated into the BlueScript vector  
25 for sequence confirmation. The predicted amino acid sequence for the precursor IL-  
18 E42A, K89A and E42A/K89A mutants are shown in SEQ ID NO:3, SEQ ID NO:4  
SEQ ID NO:5, respectively. For *E. coli* expression, each of three IL-18 DNA inserts  
were re-ligated into the pPROEX HTa vector (GIBCO/BRL) with the use of EcoRI  
and XbaI sites. The resulting protein is N-terminal fused to histidine (Fig. 1).

30

Example 3

**Protein expression and purification.** The IL18 mutant precursors were expressed in *E.coli*, affinity purified by virtue of the histidine tag and the respective mature molecules were generated by proteolytic cleavage with factor Xa.

Each of the four pPROEX HTu/IL-18 plasmids (WT and three mutants) was introduced into competent *E. coli* cells of the DHQ strain (GIBCO/BRL) and expressed as described (11). An overnight culture of 25 ml served as the inoculum for a 450 ml of LB culture containing 100 µg/ml ampicillin and grown until it reached a cell density of 0.6-1 OD<sub>600</sub>. Protein expression was induced by treatment with isopropylthiogalactoside (IPTG 0,3 mM), and incubation continued at 37°C with shaking for 3 h. The cultured bacteria cells were harvested by centrifugation (5,000 x g for 15 min at 4°C), and the pellet was suspended in 30 ml of Talon buffer (50 mM NaH<sub>2</sub>PO<sub>4</sub>/20 mM Tris-HCl/100 mM NaCl, pH 8). Cells were lysed by sonication (2 x 30 s bursts) on ice. The soluble protein was obtained by centrifugation (4,000xg for 30 min at 4°C) and applied to a 3 ml mini-Talon column (CLONTECH). The Talon column was then washed with 30 bed volumes of Talon buffer and eluted with 6 ml of 100 mM imidazole in Talon buffer. The eluant was dialyzed against factor Xa buffer (20 mM Tris-HCl/150 mM NaCl/2 mM CsCl<sub>2</sub>) at 4°C for 20 h. The 0.2 ml of Talon affinity-purified N-terminus His x 6 fusion proIL-18 was incubated with 4 µg of factor Xa enzyme (New England Biolabs) for 4 h at room temperature in the presence of 2 mM phenylmethylsulfonyl fluoride (GIBCO/BRL). The amount of IL-18 produced was monitored by a specific ELISA(R&D Systems). The amino acid sequence predicted for the mature IL-18 WT, E42A, K89A and E42A/K89A mutants are shown in sequences SEQ ID NO:6, SEQ ID NO:7 SEQ ID NO:8, respectively.

#### 25 **Example 4**

**Characterization of the E42A, K89A and E42A/K89A IL-18 mutant proteins by Western blot.** The purified IL-18 mutants were subjected to western blot analysis with a polyclonal antibody and a monoclonal antibody specific for the mature IL-18.

30 Equal amounts of talon affinity purified precursor and mature protein (after cleavage by factor Xa) the WT and mutant IL-18 forms, were resolved by SDS/PAGE

(10% acrylamide) under reducing conditions. The proteins were transferred to nitrocellulose membranes and then incubated with the primary antibodies (rabbit anti-human IL-18 polyclonal antibody or monoclonal antibody clone 8-31-4 (IgG2a) which were raised against the recombinant mature form of human IL-18 (Puren et al., 1999) which also recognize precursor IL-18). After 24 h incubation, the corresponding second antibody, goat anti-mouse or donkey anti-rabbit IgG peroxidase (Jackson Immuno Research), was added and developed by ECL (New England Nuclear Life Science Products).

The staining of proIL-18 by polyclonal rabbit anti-human IL-18 was of equal intensity for the WT and each of the three mutants. Similarly the signals obtained with the mature forms of WT and IL-18 and each of the three mutants using the polyclonal antiserum were of equal intensity. The apparent molecular weight indicated that the different IL-18 forms were of the correct size. In contrast, when the monoclonal antibody is used, the two mutants K89A and E42A/K89A, appear to stain more intensely than the WT and the E42A mutant, suggesting that the affinity of the monoclonal antibody is greater for these mutants. These results suggest that mutants K89A and E42A/K89A may have a different conformation resulting in higher affinity.

## Example 5

**Characterization of the biological activity of E42A, K89A and E42A/K89A IL-18 mutant proteins.** The purified mature forms of IL-18/ICE/Xa were analysed for the co-induction of IFN- $\gamma$  in human natural killer cells (NKO described in example 8), in PBMCs (described in example 7) and for the induction of IL-8 in PBMCs.

IL-18 does not induce IFN- $\gamma$  in these cells unless IL-12 (or IL-15) is used as a co-stimulator. Low concentrations of IL-12 (1-2 ng/ml 12 (PreproTech Rocky Hill, NJ)) induce a small amount, of IFN- $\gamma$ , however, treatment with IL-12 together with IL-18 greatly augments IFN- $\gamma$  production. IFN- $\gamma$  produced was monitored in the cell as described in example 9. The induction of IFN- $\gamma$  in NK0 cells by WT IL-18/ICE/Xa and IL-12 was found to be comparable to that induced by recombinant

mature human IL-18 resulting from ICE processing of proIL-18 (Gu et al., 1997) and IL-12. These results indicate that the IL-18 was correctly assembled in *E. coli* and correctly processed by factor Xa.

To test the activity of the mutated IL-18, the induction of IFN- $\gamma$  production by stimulation with the mutant or WT IL-18 together with IL-12 was assessed in NKO cells (statistical analyses are described in example 10). As shown in Fig. 3A, WT IL-18 was active as an inducer of IFN- $\gamma$ , beginning at 7.5 ng/ml and increasing progressively up to 60 ng/ml (the highest concentration tested). Each of the three mutated IL-18 forms exhibited biological activity greater than that of WT in these cells. For example, the single mutation E42A was twice as active as the WT form at each of the concentrations tested. The single mutation K89A was four times more active than the WT at a concentration of 7.5 ng/ml. The double mutation E2A/K89A resulted in the most active IL-18. As shown in Fig. 3A, the E42A mutated IL-18 induced 600 pg/ml IFN $\gamma$ , the maximal activity observed by pretreatment with 60 ng/ml IL-18 WT, at a concentration of 30 ng/ml, the K89A mutant at a concentration of 15 ng/ml and the double mutant at a concentration of 7.5 ng/ml. The mutants E42A, K89A and double mutant were therefore 2, 4 and 8 folds more potent than the WT, respectively.

Similar results were observed when IFN- $\gamma$  production was tested in freshly isolated human PBMCs (example 7). In these cells, the co-stimulation of IL-12 and IL-18 resulted, in IFN- $\gamma$  production, whereas neither of the two cytokines alone could induce IFN $\gamma$ . The double mutant E42A/ K89A was the most active (Fig. 3B).

The results indicate that replacement of the two charged amino acids Glu 42 and/or Lys 89 by Ala residues consistently bring about an increase in the biological activity of IL-18.

IL-18 is known to induce IL-8 in CD14<sup>+</sup> cells in PBMC preparations (described in example 7). Although IL-18 induces IL-8 production in these cells without the need of IL-12 co-stimulation, the induction of IL-8 requires higher concentrations of IL-18 than induction of IFN- $\gamma$ . Induction of IL-8 by IL-18 WT and mutant stimulation of PBMCs was therefore tested. The IL-8 produced was monitored in the cell media by the specific assay described in example 9. Figure 4

shows that although the two single mutations were comparable to the WT in the induction of IL-8, the double mutated IL-18 induced significantly more IL-8 (3.5 fold) than the wild type version.

These results indicate that the double mutant, E42A/K89A exhibits the highest biological activity.

#### Example 6

**Neutralization of IL-18 mutants by IL-18BP.** The mutations were designed in residues predicted to be important for IL-18 binding by the inhibitor IL-18BP. The ability of IL-18BP to neutralize the biological activity of IL-18, e.g IFN- $\gamma$  production (example 8), was therefore specifically assessed.

Different concentrations of IL-18BP ("a" isoform of CHO cell produced recombinant his-6-tagged human IL-18BP (supplied by Interpharm Laboratories, Ness Ziona, Israel Kim et al., 2000)) were pre-incubated with WT IL-18 or its mutated forms (30 ng/ml final concentration) and then added to cell cultures.

As shown in Fig. 4A, the 50% inhibitory concentration of IL-18BP for co-induction of IFN- $\gamma$  by WT IL-18 from NKO cells was approximately 15 ng/ml (assuming that no inhibition occurs at 3.7 ng/ml IL-18BP and this value represent 100% activity). The single mutation of E42A resulted in a similar dose-inhibitory concentration by IL-18BP.

However, when the mutant K89A was incubated with IL-18BP, its ability to act as a co-inducer of IFN- $\gamma$  in NKO cells was neutralized at a lesser extent (Fig. 5A). Only at a concentration of 120 ng/ml a statistically significant reduction in activity could be observed. In contrast, IL-18BP failed to neutralize the double IL-18 mutant E42A/K89A.

As shown in Fig. 4B IL-18 is more sensitive to neutralization by IL-18BP when tested in PBMCs rather than NKO cells. The amount of IL-18BP needed to neutralize WT IL-18 was 3.7 ng/ml, the lowest concentration tested. The single mutation E42A behaved similarly as WT IL-18, as established by the observation that low concentrations of IL18BP neutralized its biological activity in PBMCs. In contrast, the single mutation K89A was neutralized at 120 ng/ml. Similar to the



results concerning neutralization of IL-18 mutants by IL-18BP in NKO cell, the double mutant E42A/ K89A was only slightly affected by IL18-BP in PBMCs.

These results show that the mutant E89A and the double mutant E42A/K89A are less affected by the natural inhibitor IL-18BP.

5

### Example 7

**Isolation and culture of peripheral blood mononuclear cells (PBMCs) and induction of IFN- $\gamma$ .** Residual leukocytes from platelet plateletpheresis of healthy human donors were rinsed from blood tubing and subjected to centrifugation over  
10 Histopaque. PBMCs were aspirated from the interface, washed three times in pyrogen-free saline (Baxter Health Care, Mundelein, IL), and resuspended at  $5 \times 10^6$  cells per ml in RPMI 1640 medium supplemented with 10% FBS (GIBCO/BRL Grand Island, NY). The cells were cultured in flat-bottomed 96-well plates (Becton Dickinson) with RPMI 1640 medium only (control), varying concentrations of  
15 recombinant human IL-18, and WT IL-18 (ICF/Xa) or the three mutants, in the presence of 1 ng/ml IL-12. In some experiments, IL18 preparations were first mixed with polymyxin B (1  $\mu$ g/ml purchased from Sigma) before being added to the cells. Cells were incubated for 16-20 h at 37°C in humidified air with 5% CO<sub>2</sub>, and the culture supernatant was then collected for IFN- $\gamma$  measurement.

20

### Example 8

**Induction of IFN- $\gamma$  in NKO Cell Line.** The original parental NK92 cell line was obtained from Hans Klingerman (Gong et al., 1994). The human NKO cell line used in the present studies was a subclone of this cell line. NKO cells were  
25 maintained in supplemented RPMI 1640 medium containing 10% FBS and 50 pg/ml of IL-2 (R&D Systems) and 200 pg/ml of IL-15 (PeproTech). For assays, NKO cells were suspended at  $0.5 \times 10^6$  cells per ml in RPMI 1640 medium and stimulated in 0.2 ml volumes in 96-well plates with 0.5 ng/ml of IL-12 (PeproTech Rocky Hill, NJ) and different concentrations of recombinant human IL-18 WT, IL-18 (ICE/Xa), or  
30 E42A, K89A and E42A/K89A IL-18 mutants. After 16-20 h at 37°C in humidified air with 5% CO<sub>2</sub>, the culture supernatant was collected for IFN- $\gamma$  measurement.

**Example 9**

**Analysis of Cytokines.** The liquid-phase electrochemiluminescence (ECL) method was used to measure IFN- $\gamma$  (13) and IL-8 (12) in cell culture media. The amount of ECL was determined with the use of an Origen Analyzer (Igen, Gaithersburg, MD). The limit of detection of IFN- $\gamma$  and IL-8 was 62 pg/ml and 40 pg/ml, respectively.

**Example 10**

**Statistical analysis.** Data are expressed as the mean  $\pm$  SEM. Group means were compared by ANOVA, with the use of Fisher's least significant difference. Statistical significance was accepted within 95% confidence limits. ANOVA and correlation analyses were performed with the statistical packages STATVIEW 512 +(Brain Power, Calabasas, CA).

**Example 11**

**Production of the mature IL-18 mutants in CHO cells.** For expression and secretion of mature IL-18 mutants in CHO cells, the DNA sequence encoding the mature protein of wild type and mutant IL-18BP is ligated to the sequence of the DNA sequence of the signal peptide of human growth hormone (hGH) by two PCR reactions similarly to the reactions described in example 1. The template for the first PCR reaction for the amplification of each IL-18 mutant is the corresponding construct from example 2 with sense primer (Pr 9) containing overlapping sequences of IL-18 and hGH signal peptide and reverse primers (Pr 10) encoding the last 12 nucleotides of the IL-18, a stop codon and a site for a restriction enzyme. For the amplification of the growth hormone signal peptide the plasmid pXGH is used as the template with a sense primer (Pr 11) containing a site for a restriction enzyme, the first 12 nucleotides of the hGH signal peptide and the reverse primer (Pr 12), containing overlapping sequences with the hGH signal peptide and IL-18 mature protein. The templates for the second PCR performed for the amplification of the fragment encoding the signal peptide of the hGH fused to the mature sequence of the

IL-18, are the purified amplified fragments from the first PCR reaction and the primers Pr 10 and Pr 11 containing the restriction sites. The fusion fragment is purified, digested with the appropriate restriction enzymes and cloned into a mammalian expression vector.

- 5           The plasmids are used for transfecting CHO (DHFR-) cells together with a plasmid containing the mouse DHFR gene as a genetic marker. Resistant cells are isolated in a selective medium and assayed for IL-18 production by an ELISA assay.

- The stably transfected cells are subjected to several cycles of gene amplification with increasing concentrations of MTX. At the end of the gene amplification process,  
10       clones are isolated by limiting dilution. After subcloning the clone that show high specific productivity and grater stability of production is selected for production.

## REFERENCES

1. Anderson, D.M., Maraskovsky, E., Billingsley, W.L., Dougall, W.C., Tometsko, M.E., Roux, E.R., Teepe, M.C., DuBose, R.F., Cosman, D., Galibert, L. (1997) "A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function." *Nature*, 390, 175-179.
2. Bazan, J. F., Timans, J. C. and Kaselein, R. A. (1996) "A newly defined interleukin-1?" *Nature* 379, 591.
3. Born, T.L., Morrison, L.A., Esteban, D.J., VandenBos, T., Thebeau, L.G., Chen, N., Spriggs, M.K., Sims, J.E., Buller, R.M. (2000) "A poxvirus protein that binds to and inactivates IL-18, and inhibits NK cell response." *J Immunol* 164,3246-54.
4. Childs, R., Chernoff, A., Contentin, N., Bahceci, E., Schrupp, D., Leitman, S., Read, E.J., Tisdale, J., Dunbar, C., Linehan, W.M., Young, N.S., Barrett, A.J. (2000) "Regression of metastatic renal-cell carcinoma after nonmyeloablative allogeneic peripheral-blood stem-cell transplantation." *N Engl J Med* 343, 750-8.
5. Cho, D., Kim, T.G., Lee, W., Hwang, Y.I., Cho, H.I., Han, H., Kwon, O., Kim, D., Park, H., Houh, D. (2000) "Interleukin-18 and the costimulatory molecule B7-1 have a synergistic anti-tumor effect on murine melanoma; implication of combined immunotherapy for poorly immunogenic malignancy" *J Invest Dermatol* ,114, 928-34.
6. Coughlin, C.M., Salhany, K.E., Wysocka, M., Aruga, E., Kurzawa, H., Chang, A.E., Hunter, C.A., Fox, J.C., Trinchieri, G. and Lee, W.M. (1998) "Interleukin-12 and interleukin-18 synergistically induce murine tumor regression which involves inhibition of angiogenesis." *J Clin Invest* Mar, 101,1441-52.
7. Engelmann, H., Aderka, D., Rubinstein, M., Rotman, D. and Wallach. D. (1989)" A tumor necrosis factor-binding protein purified to homogeneity from human urine protects cells from tumor necrosis factor toxicity" *J. Biol. Chem.* 264,11974-11980.
8. Engelmann, H., Novick, D. and Wallach, D. (1990)"Two tumor necrosis factor-binding proteins purified from human urine. Evidence for immunological cross-reactivity with cell surface tumor necrosis factor receptors." *J. Biol. Chem.* 265,1531-1536.

9. Ghayur, T., Banerjee, S., Hugunin, M., Butler, D., Herzog, L., Carter, A., Quintal, L., Sekut, L., Talanian, R., Paskind, M., Wong, W., Kamen, R., Tracey, D., and Allen, H. (1997) "Caspase-1 processes IFN-gamma-inducing factor and regulates LPS-induced IFN-gamma production." *Nature* 386, 619-623.
- 5 10. Gollob, J.A., Mier, J.W., Veenstra, K., McDermott, D.F., Clancy, D., Clancy, M., Atkins, M.B. (2000) "Phase I trial of twice-weekly intravenous interleukin 12 in patients with metastatic renal cell cancer or malignant melanoma: ability to maintain IFN-gamma induction is associated with clinical response." *Clin Cancer Res* , 5, 1678-92.
- 10 11. Gong, J. H., Maki, G., Klingemann, H. G. (1994) "Characterization of a human cell line (NK-92) with phenotypical and functional characteristics of activated natural killer cells." *Leukemia* 8:652.
12. Gu, Y., Kuida, K., Tsutsui, H., Ku, G., Hsiao, K., Fleming, M. A., Hayashi, N., Higashino, K., Okamura, H., Nakanishi, K., Kurimoto, M., Tanimoto, T., Flavell, R. A., Sato, V., Harding, M. W., Livingston, D. J., and Su, M. S. (1997) "Activation of interferon-gamma inducing factor mediated by interleukin-1beta converting enzyme." *Science* 275, 206-209
- 15 13. Kim, S.H., Eisenstein, M., Reznikov, L., Fantuzzi, G., Novick, D., Rubinstein, M. and Dinarello, C.A.(2000)" Structural requirements of six naturally occurring isoforms of the IL-18 binding protein to inhibit IL-18." *Proc Natl Acad Sci U S A*, 97, 1190-5.
- 20 14. Kohno, K., J. Kataoka, T. Ohtsuki, Y. Suemoto, I. Okamoto, M. Usui, M. Ikeda, and M. Kurimoto. (1997) "IFN-gamma-inducing factor (IGIF) is a costimulatory factor on the activation of Th1 but not Th2 cells and exerts its effect independently of IL-12." *J. Immunol.* 158:1541-1550.
- 25 15. Kugler, A., Stuhler, G., Walden, P., Zoller, G., Zobywalski, A., Brossart, P., Trefzer, U., Ullrich, S., Muller, C.A., Becker, V., Gross, A.J., Hemmerlein, B., Kanz, L., Muller, G.A., Ringert, R.H.( 2000) " Regression of human metastatic renal cell carcinoma after vaccination with tumor cell-dendritic cell hybrids. " *Nat Med*,3, 332-6.

16. Nakamura, K., Okamura, H., Wada, M., Nagata, K. and Tamura, T. (1989).  
 "Endotoxin-induced serum factor that stimulates gamma interferon production."  
*Infect. Immun.*, 57,590-5 issn: 0019-9567.
17. Nakamura, K., Okamura, H., Nagata, K., Komatsu, T. and Tamura, T. (1993)  
 5 "Purification of a factor which provides a costimulatory signal for gamma interferon  
 production." *Infect. Immun.*, 61, 64-70
18. Novick, D., Engelmann, H., Wallach, D. and Rubinstein. M. (1989) "Soluble  
 cytokine receptors are present in normal human urine." *J. Exp. Med.*, 170,1409-14.
19. Novick, D., Cohen, B. and Rubinstein, M. (1992) "Soluble Interferon-alpha  
 10 Receptor Molecules Are Present in Body Fluids." *FEBS Lett*, 314,445-8.
20. Novick, D., Cohen, B. and Rubinstein, M. (1994) "The Human Interferon  
 alpha/beta Receptor - Characterization and Molecular Cloning." *Cell* 77,391-400.
21. Novick, D., Kim,S., Fantuzzi, G., Reznikov, L.L., Dinarello, C.A. and Rubinstein,  
 M. (1999) "Interleukin-18 Binding Protein: A Novel Modulator of the Th1  
 15 Cytokine Response. *Immunity*, 10, 127,36.
22. Okamura, H., Tsutsui, H., Komatsu, T., Yutsudo, M., Hakura, A., Tanimoto, T.,  
 Torigoe, K., Okura, T., Nukada, Y., Hattori, K., Akita, K., Namba, M., Tanabe, F.,  
 Konishi, K., Fukuda, S., and Kurimoto, M. (1995) "Cloning of a new cytokine that  
 induces IFN-gamma production by T cells." *Nature*, 378, 88-91
- 20 23. Puren, A.J., Fantuzzi, G., Dinarello, C.A. (1999) "Gene expression, synthesis, and  
 secretion of interleukin 18 and interleukin 1beta are differentially regulated in human  
 blood mononuclear cells and mouse spleen cells." *Proc Natl Acad Sci U S A*,96,2256-  
 61.
24. Seki, .S, Habu, Y., Kawamura, T., Takeda, K., Dobashi, H., Ohkawa, T.,  
 25 Hiraide, H. (2000) " The liver as a crucial organ in the first line of host  
 defense: the roles of Kupffer cells, natural killer (NK) cells and NK1.1 Ag+  
 T cells in T helper 1 immune responses." *Immunol Rev*, 174,35-46.
25. Simonet, W.S., Lacey, D.L., Dunstan, C.R., Kelley, M., Chang, M.S.,  
 Luthy, R., Nguyen, H.Q., Wooden, S., Bennett, L., Boone, T., Shimamoto,  
 30 G., DeRose, M., Elliott, R., Colombero, A., Tan, H.L., Trail, G., Sullivan,  
 J., Davy, E., Bucay, N., Renshaw-Gegg, L., Hughes, T.M., Hill, D.,

- Pattison, W., Campbell, P., Boyle, W.J. (1997). "Osteoprotegerin: a novel secreted protein involved in the regulation of bone density." *Cell*, 89, 309-19.
26. Slavin, S. (2000) "Immunotherapy of cancer with alloreactive lymphocytes." *N Engl J Med*, 343,802-3.
27. Slavin, S., Or, R., Prighozina, T., Gurevitch, O., Aker, M., Panighari, S., Shapira, M., Nagle, A. (2000) "Immunotherapy of hematologic malignancies and metastatic solid tumors in experimental animals and man" *Bone Marrow Transplant Suppl*, 2:S54-7.
28. Tsutsui, H., K. Nakanishi, K. Matsui, K. Higashino, H. Okamura, Y. Miyazawa, and K. Kaneda. (1996) "IFN-gamma-inducing factor up-regulates Fas ligand-mediated cytotoxic activity of murine natural killer cell clones." *J. Immunol.*, 157,3967-73 issn: 0022-1767.
29. Tuting, T., Wilson, C.C., Martin, D.M., Kasamon, Y.L., Rowles, J., Ma, D.I., Slingluff, C.L., Wagner, S.N., van der Bruggen, P., Baar, J., Lotze, M.T., Storkus, W.J. (1998) "Autologous human monocyte-derived dendritic cells genetically modified to express melanoma antigens elicit primary cytotoxic T cell responses in vitro: enhancement by cotransfection of genes encoding the Th1-biasing cytokines IL-12 and IFN-alpha." *J Immunol*, 160, 1139-47.
30. Urushihara, N., Iwagaki, H., Yagi, T., Kohka, H., Kobashi, K., Morimoto, Y., Yoshino, T., Tanimoto, T., Kurimoto, M., Tanaka, N. (2000) "Elevation of serum interleukin-18 levels and activation of Kupffer cells in biliary atresia." *J Pediatr Surg*, 35,446-9.
31. Ushio, S., Namba, M., Okura, T., Hattori, K., Nukada, Y., Akita, K., Tanabe, F., Konishi, K., Micallef, M., Fujii, M., Torigoe, K., Tanimoto, T., Fukuda, S., Ikeda, M., Okamura, H., and Kurimoto, M. (1996) *J. Immunol.*, 156, 4274-9.
32. Vigers, G.P., Anderson, L.J., Caffes, P., Brandhuber, B.J. (1997) "Crystal structure of the type-I interleukin-1 receptor complexed with interleukin-1 beta." *Nature*, 386,190-4.

33. Xiang, Y. and Moss, B.( 1999)" IL-18 binding and inhibition of interferon gamma induction by human poxvirus-encoded proteins." *Proc Natl Acad Sci U S A*, 96,11537-42.

34. Yasuda, H., Shima, N., Nakagawa, N., Mochizuki, S.I., Yano, K., Fujise, N., Sato,  
5 Y., Goto, M., Yamaguchi, K., Kuriyama, M., Kanno, T., Murakami, A., Tsuda, E.,  
Morinaga, T., Higashio, K. (1998) " Identity of osteoclastogenesis inhibitory  
factor (OCIF) and osteoprotegerin (OPG): a mechanism by which OPG/OCIF  
inhibits osteoclastogenesis in vitro." *Endocrinology*, 139, 1329-37.

10 The references cited above and throughout the specification are herein  
incorporated by reference in their entirety.



**CLAIMS**

We claim:

1. An IL-18 mutant polypeptide, comprising mutations in one or more amino acid residues which are involved in its interaction with IL-18 binding protein.
- 5 2. The polypeptide according to claim 1, wherein the mutations are substitutions, additions or deletions.
3. The polypeptide according to claim 2, wherein the substitutions are non-conservative.
4. The polypeptide according to anyone of claims 1, 2 or 3, wherein the mutation  
10 is in a residue selected from a group consisting of Glu-42, Ile-85, Met-87, Lys-89, Met-96, Asp-130, Lys-132, Pro-143, Met-149, and Leu-189.
5. The polypeptide according to claim 4, wherein the mutation is in a residue selected from the group consisting of Glu-42 and Lys-89.
6. The polypeptide according to claim 5, wherein the mutation is in a Glu-42  
15 residue.
7. The polypeptide according to claim 5, wherein the mutation is in a Lys-89 residue.
8. The polypeptide according to claim 5, wherein the mutation is in a Glu-42 residue and a Lys-89 residue.
- 20 9. The polypeptide according to claim 6, wherein the Glu-42 residue is replaced with a non-polar amino acid.
10. The polypeptide according to claim 9, wherein the Glu-42 is replaced with an Ala residue.
11. The polypeptide according to claim 7, wherein the Lys-89 residue is replaced  
25 with a non-polar amino acid.
12. The polypeptide according to claim 11, wherein the Lys-89 is replaced with an Ala residue.
13. The polypeptide according to claim 8, wherein both the Glu-42 residue and the Lys-89 residues are replaced with a non-polar amino acid.
- 30 14. The polypeptide according to claim 13, where both the Glu-42 and the Lys-89 residues are replaced with an Ala residue.

15. An isolated DNA encoding a polypeptide according to any one of claims 1-14.
16. The DNA according to claim 15, wherein the polypeptide has amino acid sequence of SEQ ID NO:3.
17. The DNA according to claim 15, wherein the polypeptide has amino acid sequence of encoding the polypeptide of SEQ ID NO:4.
18. The DNA according to claim 15, wherein the polypeptide has amino acid sequence of encoding the polypeptide of SEQ ID NO:5.
19. The DNA according to claim 15, wherein the polypeptide has amino acid sequence of encoding the polypeptide of SEQ ID NO:6.
20. The DNA according to claim 15, wherein the polypeptide has amino acid sequence of encoding the polypeptide of SEQ ID NO:7.
21. The DNA according to claim 15, wherein the polypeptide has amino acid sequence of encoding the polypeptide of SEQ ID NO:8.
22. A DNA according to any one of claims 15, 19, 20 and 21 further comprising a nucleic acid sequence encoding a signal peptide.
23. The DNA according to claim 22, wherein the signal peptide is that of a growth hormone.
24. A vector comprising a DNA according to any one of claims 15 to 23, wherein said vector is capable of expressing the polypeptide encoded by said DNA in an appropriate host cell.
25. The vector according to claim 24, wherein the host cell is prokaryotic.
26. The vector according to claim 25, wherein the DNA encodes a polypeptide selected from a group consisting of SEQ ID NO:3, SEQ ID NO:4 and SEQ ID NO:5.
27. The vector according to claim 24, wherein the host cell is an eukaryotic cell.
28. The vector according to claim 27, wherein the DNA encodes a polypeptide selected from a group consisting of SEQ ID NO:6, SEQ ID NO:7 and SEQ ID NO:8.
29. The vector according to claim 27, comprising DNA according to claim 22 or 23.

30. The vector according to claim 28, wherein said DNA is ligated to the sequence encoding the human growth hormone signal peptide.
31. A pharmaceutical composition comprising a polypeptide according to any one of claims 1 to 14 and a pharmaceutically acceptable carrier for the treatment of diseases which are prevented or alleviated by Th1 responses.
32. The pharmaceutical composition according to claim 31, for the treatment of cancer.
33. The pharmaceutical composition according to claim 31, for the treatment of viral diseases.




Figure 1 A

Tyr	Phe	Gly	Lys	Leu	<b>Glu</b>	Ser	Lys	Leu	Ser	Val	Ile	Arg	Asn	Leu	Asn
1				5					10					15	
<b>Asp</b>	<b>Gln</b>	<b>Val</b>	<b>Leu</b>	<b>Phe</b>	<b>Ile</b>	<b>Asp</b>	<b>Gln</b>	<b>Gly</b>	<b>Asn</b>	<b>Arg</b>	<b>Pro</b>	<b>Leu</b>	<b>Phe</b>	<b>Glu</b>	<b>Asp</b>
		20						25					30		
<b>Met</b>	<b>Thr</b>	<b>Asp</b>	Ser	Asp	Cys	Arg	Asp	Asn	Ala	Pro	Arg	Thr	Ile	Phe	Ile
		35					40					45			
Ile	Ser	Met	<b>Tyr</b>	<b>Lys</b>	<b>Asp</b>	<b>Ser</b>	Gln	Pro	Arg	Gly	Met	Ala	Val	Thr	Ile
	50					55					60				
Ser	Val	Lys	Cys	Glu	Lys	Ile	Ser	Thr	Leu	Ser	Cys	Glu	Asn	Lys	Ile
65					70					75				80	
Ile	Ser	Phe	Lys	Glu	Met	Asn	Pro	Pro	Asp	Asn	Ile	Lys	Asp	Thr	Lys
			85						90					95	
Ser	Asp	Ile	Ile	Phe	Phe	Gln	Arg	Ser	Val	Pro	Gly	His	Asp	Asn	Lys
			100					105					110		
Met	Gln	Phe	Glu	Ser	Ser	Ser	Tyr	Glu	Gly	Tyr	Phe	Leu	Ala	Cys	Glu
		115					120					125			
Lys	Glu	Arg	Asp	Leu	Phe	Lys	Leu	Ile	Leu	Lys	Lys	Glu	Asp	Glu	Leu
	130					135						140			
Gly	Asp	Arg	Ser	Ile	Met	Phe	Thr	Val	Gln	Asn	Glu	Asp			
145					150					155					

SEQ ID NO: 2

Figure 1 B

**Figure 2**






Figure 3

**Figure 4**



**Figure 5**

## SEQUENCE LISTING

<110> Ares Trading S.A  
Dinareello, Charles  
Kim, Soo Hyun

<120> InterLeukin-18 Mutants, Their Production and use

<130> 475

<150> 60/274,327

<151> 2001-03-08

<160> 18

<170> PatentIn version 3.1

<210> 1

<211> 582

<212> DNA

<213> Homo sapiens

<400> 1

atggctgctg aaccagtaga agacaattgc atcaactttg tggcaatgaa atttattgac  
60

aatacgcttt actttatagc tgaagatgat gaaaacctgg aatcagatta ctttggcaag  
120

cttgaatcta aattatcagt cataagaaat ttgaatgacc aagttctctt cattgaccaa  
180

ggaaatcggc ctctatttga agatatgact gattctgact gtagagataa tgcaccccg  
240

accatattta ttataagtat gtataaagat agccagccta gaggtatggc tgtaactatc  
300

tctgtgaagt gtgagaaaat ttcaactctc tcctgtgaga acaaaattat ttcctttaag  
360

gaaatgaatc ctctgataa catcaaggat acaaaaagtg acatcatatt ctttcagaga  
420

agtgtcccag gacatgataa taagatgcaa tttgaatctt catcatacga aggatacttt  
480

ctagcttggtg aaaaagagag agaccttttt aaactcattt tgaaaaaaga ggatgaattg  
540

ggggatagat ctataatggt cactgttcaa aacgaagact ag  
582

<210> 2  
 <211> 157  
 <212> PRT  
 <213> Homo sapiens

<400> 2

Tyr Phe Gly Lys Leu Glu Ser Lys Leu Ser Val Ile Arg Asn Leu Asn  
 1 5 10 15

Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro Leu Phe Glu Asp  
 20 25 30

Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg Thr Ile Phe Ile  
 35 40 45

Ile Ser Met Tyr Lys Asp Ser Gln Pro Arg Gly Met Ala Val Thr Ile  
 50 55 60

Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys Glu Asn Lys Ile  
 65 70 75 80

Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile Lys Asp Thr Lys  
 85 90 95

Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asn Lys  
 100 105 110

Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe Leu Ala Cys Glu  
 115 120 125

Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys Glu Asp Glu Leu  
 130 135 140

Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu Asp  
 145 150 155

<210> 3  
 <211> 193  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Synthetic PRT Sequence

&lt;400&gt; 3

Met Ala Ala Glu Pro Val Glu Asp Asn Cys Ile Asn Phe Val Ala Met  
 1 5 10 15

Lys Phe Ile Asp Asn Thr Leu Tyr Phe Ile Ala Glu Asp Asp Glu Asn  
 20 25 30

Ile Glu Gly Arg Tyr Phe Gly Lys Leu Ala Ser Lys Leu Ser Val Ile  
 35 40 45

Arg Asn Leu Asn Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro  
 50 55 60

Leu Phe Glu Asp Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg  
 65 70 75 80

Thr Ile Phe Ile Ile Ser Met Tyr Lys Asp Ser Gln Pro Arg Gly Met  
 85 90 95

Ala Val Thr Ile Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys  
 100 105 110

Glu Asn Lys Ile Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile  
 115 120 125

Lys Asp Thr Lys Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly  
 130 135 140

His Asp Asn Lys Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe  
 145 150 155 160

Leu Ala Cys Glu Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys  
 165 170 175

Glu Asp Glu Leu Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu  
 180 185 190

Asp

<210> 4  
 <211> 193  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Synthetic PRT Sequence

<400> 4

Met Ala Ala Glu Pro Val Glu Asp Asn Cys Ile Asn Phe Val Ala Met  
 1 5 10 15

Lys Phe Ile Asp Asn Thr Leu Tyr Phe Ile Ala Glu Asp Asp Glu Asn  
 20 25 30

Ile Glu Gly Arg Tyr Phe Gly Lys Leu Glu Ser Lys Leu Ser Val Ile  
 35 40 45

Arg Asn Leu Asn Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro  
 50 55 60

Leu Phe Glu Asp Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg  
 65 70 75 80

Thr Ile Phe Ile Ile Ser Met Tyr Ala Asp Ser Gln Pro Arg Gly Met  
 85 90 95

Ala Val Thr Ile Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys  
 100 105 110

Glu Asn Lys Ile Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile  
 115 120 125

Lys Asp Thr Lys Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly  
 130 135 140

His Asp Asn Lys Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe  
 145 150 155 160

Leu Ala Cys Glu Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys  
 165 170 175

Glu Asp Glu Leu Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu  
 180 185 190

Asp

<210> 5  
 <211> 193  
 <212> PRT  
 <213> Artificial Sequence  
 <220>  
 <223> Synthetic PRT Sequence  
 <400> 5

Met Ala Ala Glu Pro Val Glu Asp Asn Cys Ile Asn Phe Val Ala Met  
 1 5 10 15

Lys Phe Ile Asp Asn Thr Leu Tyr Phe Ile Ala Glu Asp Asp Glu Asn  
 20 25 30

Ile Glu Gly Arg Tyr Phe Gly Lys Leu Ala Ser Lys Leu Ser Val Ile  
 35 40 45

Arg Asn Leu Asn Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro  
 50 55 60

Leu Phe Glu Asp Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg  
 65 70 75 80

Thr Ile Phe Ile Ile Ser Met Tyr Ala Asp Ser Gln Pro Arg Gly Met  
 85 90 95

Ala Val Thr Ile Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys  
 100 105 110

Glu Asn Lys Ile Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile  
 115 120 125

Lys Asp Thr Lys Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly  
 130 135 140

His Asp Asn Lys Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe  
145 150 155 160

Leu Ala Cys Glu Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys  
165 170 175

Glu Asp Glu Leu Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu  
180 185 190

Asp

<210> 6

<211> 157

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic PRT Sequence

<400> 6

Tyr Phe Gly Lys Leu Ala Ser Lys Leu Ser Val Ile Arg Asn Leu Asn  
1 5 10 15

Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro Leu Phe Glu Asp  
20 25 30

Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg Thr Ile Phe Ile  
35 40 45

Ile Ser Met Tyr Lys Asp Ser Gln Pro Arg Gly Met Ala Val Thr Ile  
50 55 60

Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys Glu Asn Lys Ile  
65 70 75 80

Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile Lys Asp Thr Lys  
85 90 95

Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asn Lys  
100 105 110

Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe Leu Ala Cys Glu  
115 120 125

Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys Glu Asp Glu Leu  
130 135 140

Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu Asp  
145 150 155

<210> 7  
<211> 157  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Synthetic PRT Sequence

<400> 7

Tyr Phe Gly Lys Leu Glu Ser Lys Leu Ser Val Ile Arg Asn Leu Asn  
1 5 10 15

Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro Leu Phe Glu Asp  
20 25 30

Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg Thr Ile Phe Ile  
35 40 45

Ile Ser Met Tyr Ala Asp Ser Gln Pro Arg Gly Met Ala Val Thr Ile  
50 55 60

Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys Glu Asn Lys Ile  
65 70 75 80

Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile Lys Asp Thr Lys  
85 90 95

Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asn Lys  
100 105 110

Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe Leu Ala Cys Glu  
115 120 125



Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys Glu Asp Glu Leu  
 130 135 140

Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu Asp  
 145 150 155

<210> 8

<211> 157

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic PRT Sequence

<400> 8

Tyr Phe Gly Lys Leu Ala Ser Lys Leu Ser Val Ile Arg Asn Leu Asn  
 1 5 10 15

Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro Leu Phe Glu Asp  
 20 25 30

Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg Thr Ile Phe Ile  
 35 40 45

Ile Ser Met Tyr Ala Asp Ser Gln Pro Arg Gly Met Ala Val Thr Ile  
 50 55 60

Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys Glu Asn Lys Ile  
 65 70 75 80

Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile Lys Asp Thr Lys  
 85 90 95

Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asn Lys  
 100 105 110

Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe Leu Ala Cys Glu  
 115 120 125

Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys Glu Asp Glu Leu  
 130 135 140

Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu Asp  
 145 150 155

<210> 9  
 <211> 193  
 <212> PRT  
 <213> Artificial Sequence  
  
 <220>  
 <223> Synthetic PRT Sequence  
  
 <400> 9

Met Ala Ala Glu Pro Val Glu Asp Asn Cys Ile Asn Phe Val Ala Met  
 1 5 10 15

Lys Phe Ile Asp Asn Thr Leu Tyr Phe Ile Ala Glu Asp Asp Glu Asn  
 20 25 30

Ile Glu Gly Arg Tyr Phe Gly Lys Leu Lys Ser Lys Leu Ser Val Ile  
 35 40 45

Arg Asn Leu Asn Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro  
 50 55 60

Leu Phe Glu Asp Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg  
 65 70 75 80

Thr Ile Phe Ile Ile Ser Met Tyr Lys Asp Ser Gln Pro Arg Gly Met  
 85 90 95

Ala Val Thr Ile Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys  
 100 105 110

Glu Asn Lys Ile Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile  
 115 120 125

Lys Asp Thr Lys Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly  
 130 135 140

His Asp Asn Lys Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe  
 145 150 155 160

Leu Ala Cys Glu Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys  
                     165                    170                    175

Glu Asp Glu Leu Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu  
                     180                    185                    190

Asp

<210> 10  
 <211> 157  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Synthetic PRT Sequence

<400> 10

Tyr Phe Gly Lys Leu Lys Ser Lys Leu Ser Val Ile Arg Asn Leu Asn  
 1                    5                    10                    15

Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro Leu Phe Glu Asp  
                     20                    25                    30

Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg Thr Ile Phe Ile  
                     35                    40                    45

Ile Ser Met Tyr Lys Asp Ser Gln Pro Arg Gly Met Ala Val Thr Ile  
                     50                    55                    60

Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys Glu Asn Lys Ile  
 65                    70                    75                    80

Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile Lys Asp Thr Lys  
                     85                    90                    95

Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asn Lys  
                     100                    105                    110

Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe Leu Ala Cys Glu  
                     115                    120                    125

Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys Glu Asp Glu Leu  
130 135 140

Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu Asp  
145 150 155

<210> 11  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Synthetic DNA Sequence

<400> 11  
atatgaattc atggctgctg aaccagtag  
29

<210> 12  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Synthetic DNA Sequence

<400> 12  
aaagtaacgt ccttcgatgt tttc  
24

<210> 13  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Synthetic DNA Sequence

<400> 13  
gaaaacatcg aaggacgtta cttt  
24

<210> 14  
<211> 31  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Synthetic DNA Sequence

<400> 14

atatggatcc tagtcttcgt tttgaacagt g  
31

<210> 15

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic DNA Sequence

<400> 15

taatttagat gcaagcttgc c  
21

<210> 16

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic DNA Sequence

<400> 16

ggcaagcttg catctaaatt a  
21

<210> 17

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic DNA Sequence

<400> 17

ctggctatct gcatacatat t  
21

<210> 18

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic DNA Sequence

<400> 18  
agtatgtatg cagatagcca g  
21

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
19 December 2002 (19.12.2002)

PCT

(10) International Publication Number  
**WO 02/101049 A3**

(51) International Patent Classification<sup>7</sup>: **C12N 15/24**,  
C07K 14/54, A61K 38/20, A61P 31/12, 35/00

(21) International Application Number: PCT/IB02/02344

(22) International Filing Date: 8 March 2002 (08.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/274,327 8 March 2001 (08.03.2001) US

(71) Applicant: **ARES TRADING S.A.** [CH/CH]; Le  
Château, CH-2028 Vaumarcus (CH).

(72) Inventors: **DINARELLO, Charles**; 333 15th Street,  
Boulder, CO 80302 (US). **KIM, Soo-Hyun**; 19833 E.  
Villanova, Aurora, CO 80013 (US).

(74) Agent: **SERONO INTERNATIONAL S.A.**; Intellectual  
Property, 12, chemin des Aulx, CH-1228 Plan-les-Ouates  
(CH).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,  
YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent  
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,  
NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:  
18 December 2003

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **INTERLEUKIN-18 MUTANTS, THEIR PRODUCTION AND USE**

(57) Abstract: The invention provides mutants of IL-18 with lower affinity to IL-18BP than the wild type IL-18 molecule.

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/IB 02/02344

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/24 C07K14/54 A61K38/20 A61P31/12 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, EPO-Internal, WPI Data, PAJ, MEDLINE, EMBL, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	KIM SOO-HYUN M ET AL: "Site-specific mutations in the mature form of human IL-18 with enhanced biological activity and decreased neutralization by IL-18 binding protein." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES, vol. 98, no. 6, 13 March 2001 (2001-03-13), pages 3304-3309, XP002226879 March 13, 2001 ISSN: 0027-8424 the whole document	1-33
Y	EP 0 845 530 A (HAYASHIBARA BIOCHEM LAB) 3 June 1998 (1998-06-03) cited in the application the whole document	1-33
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

13 January 2003

Date of mailing of the international search report

31/01/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Bucka, A



## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>KIM S-H ET AL: "STRUCTURAL REQUIREMENTS OF SIX NATURALLY OCCURRING ISOFORMS OF THE IL-18 BINDING PROTEIN TO INHIBIT IL-18" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, US, vol. 97, no. 3, 1 February 2000 (2000-02-01), pages 1190-1195, XP000941910 ISSN: 0027-8424 cited in the application page 1193, right-hand column; figure 6</p>	1-33
Y	<p>RUBINSTEIN M ET AL.: "Cytokines, cytokine receptors and cytokine inhibitors" LIFE SCIENCE BOOK 2000, 'Online! 2000, XP002226976 WEIZMANN INSTITUTE OF SCIENCE, REHOVOT, ISRAEL Retrieved from the Internet: &lt;URL:http://www.weizmann.ac.il/Biology/open-day/images/rubinm.pdf&gt; 'retrieved on 2003-01-13! page 98 -page 99; figure 2 &amp; "Life Science Book 2000" 2000, WEIZMANN INSTITUTE OF SCIENCE, REHOVOT, ISRAEL</p>	1-33
A	<p>VIGERS GUY P A ET AL: "Crystal structure of the type-I interleukin-1 receptor complexed with interleukin-1-beta." NATURE (LONDON), vol. 386, no. 6621, 1997, pages 190-194, XP001109468 ISSN: 0028-0836 cited in the application the whole document</p>	1-33
A	<p>DINARELLO C A ET AL: "OVERVIEW OF INTERLEUKIN-18: MORE THEN AN INTERFERON-GAMMA INDUCING FACTOR" JOURNAL OF LEUKOCYTE BIOLOGY, FEDERATION OF AMERICAN SOCIETIES FOR EXPERIMENTAL, US, vol. 63, no. 6, June 1998 (1998-06), pages 658-664, XP000973451 ISSN: 0741-5400 the whole document</p>	1-33

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
EP 0845530	A	03-06-1998	JP	10262686 A	06-10-1998
			EP	0845530 A2	03-06-1998
			US	6476197 B1	05-11-2002
<hr/>					